

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The role of hepatic enzymes in Crohn's disease

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1632784> since 2018-10-31T17:20:28Z

Published version:

DOI:10.1007/s00384-017-2829-1

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 Davide Giuseppe Ribaldone,¹ Marco Astegiano,¹ Rinaldo Pellicano²

2
3 The role of hepatic enzymes in Crohn's disease

4
5 Department(s) and institution(s)

6 ¹MD, Gastroenterology - U, General and Specialist Medicine Department, Città della
7 Salute e della Scienza of Turin, C.so Bramante 88, 10126 Turin, Italy, ²MD,
8 Department of Gastroenterology, Molinette Hospital, Turin, Italy

9
10 Corresponding Author: Dr. Davide Giuseppe Ribaldone, General and Specialist
11 Medicine Department, Città della Salute e della Scienza of Turin, C.so Bramante 88,
12 10126 Turin, tel 00390116335208, fax 00390116336752, Italy. E-mail:
13 davrib_1998@yahoo.com

14
15 Davide Giuseppe Ribaldone ORCID iD 0000-0002-9421-3087

16 Marco Astegiano ORCID iD 0000-0003-0916-1188

17 Rinaldo Pellicano ORCID iD 0000-0003-3438-0649

32 Abstract

33 N/A

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64 Keywords: biopsy, Crohn disease, elastography, histology, inflammatory bowel
65 diseases, liver

Sir,

a systematic review has reported that, in Crohns' disease (CD) patients, the prevalence of cholelithiasis ranged from 11% to 34%, that of primary sclerosing cholangitis (PSC) from 1.2% to 3.4%, that of fatty liver disease 23% and hepatic amyloidosis occurred in <1% [1]. In a prospective, single-blind study we evaluated the prevalence of histological changes in the liver of patients with CD, without alterations of both liver biochemical tests and ultrasound, and their prognostic significance. The patients underwent liver biopsy at the time of intestinal resection. Exclusion criteria were a known liver disease. Thereafter, patients were clinically monitored every 6 months, upper abdomen ultrasound was performed at least every 12 months. Finally, after a mean interval of 14 years from liver biopsy, these patients were assessed using the Fibroscan® (Echosens®, Paris, France). Ultrasound examination in the pre-operative step showed steatosis in 10 (29%) patients. At biopsy specimens alterations in 60% of patients, without serious liver injuries, were found. No evidence of a significant liver damage progression after a mean period of 14 years were found. The average result (5.2 ± 1.2 kPa) obtained performing Fibroscan® was comparable to that (5.30 ± 1.45 kPa, $p = 0.63$) reported in healthy subjects [2].

A recent interesting retrospective study was performed in 383 CD patients newly diagnosed (not treated). One patient with chronic liver disease (small duct PSC) was excluded. Of the 383 patients included in this study, 131 had liver test abnormalities (34.1%), but liver diseases were not found, apart from liver steatosis in 6% of patients [3] (*versus* 29% in the previous study [2]).

The two studies [2, 3] agree that, considering the cost/benefit ratio, patients with CD should be considered as healthy from the liver perspective, without the need for additional biochemical and instrumental examinations than the general population, unless the presence of clinical or biochemical suspicion of liver disease.

In the more recent study [3], however, the authors found that, patients with liver test abnormalities, without an hepatic disease, more often developed complicated CD behaviour and more often needed hospitalization or surgery within 5 years of diagnosis than patients without liver test abnormalities. Patients with a C-reactive protein (CRP) <16 mg/L but with liver test abnormalities had a higher risk of developing complicated disease compared to those without liver test abnormalities. This demonstrates that the presence of liver test abnormalities does not merely reflect a higher CRP concentration, but may be a more sensitive indicator of an increased risk of complicated disease behaviour than CRP. At multivariate analysis, the presence of liver test abnormalities was independent risk factors for complicated disease behaviour; additionally, the presence of liver test abnormalities

95 was independently associated with an increased risk of hospitalizations (HR 1.7, $p = 0.023$) as well as surgery
96 (HR 2.3, $p = 0.015$) [3].

97 In conclusion, in the absence of a known liver disease or of a risk factor for hepatic injury (e.g., a potentially
98 hepatotoxic drugs), liver enzymes in CD do not need to be routinely measured. However, when increased, liver
99 enzymes could predict a more aggressive CD behaviour.

100

101 **Compliance with ethical standards**

102

103 **Funding** None to declare.

104

105 **Conflict of Interest** None to declare.

106

107 **Ethical approval** The study was conducted in accordance with ICH Good Clinical Practice
108 guidelines, the Declaration of Helsinki, and local laws and regulations.

109

110 **Informed consent** Informed consent due to the observational study has been obtained in the
111 cited studies.

112

113

114

115

116

117

118

119

References

1. Gizard E, Ford AC, Bronowicki JP, Peyrin-Biroulet L (2014) Systematic review: The epidemiology of the hepatobiliary manifestations in patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 40:3–15.
2. Ribaldone DG, Garavagno M, Pellicano R, Bresso F, Fagoonee S, David E, Sapone N, Bonagura AG, Resegotti A, Astegiano M (2015) Prevalence and prognostic value of hepatic histological alterations in patients with Crohn's disease. *Scand J Gastroenterol* 50:1463-1468.
3. Barendregt J, de Jong M, Haans JJ, van Hoek B, Hardwick J, Veenendaal R, van der Meulen A, Srivastava N, Stuyt R, Maljaars J (2017) Liver test abnormalities predict complicated disease behaviour in patients with newly diagnosed Crohn's disease. *Int J Colorectal Dis* 32:459-467.